



TM

Concentrate\*

**Selective broadleaf weed herbicide**  
*Specially formulated for control of broadleaf weeds in turf<sup>^</sup>(parks, lawns, golf courses, cemeteries, sports fields, and other grassy areas.)*

**Kills over 60 listed broadleaf weeds without harming lawns<sup>^</sup>**

**Low-Odor Formulation**

**Microtechnology<sup>‡</sup> improves penetration of reduced active ingredient formulation.**

\*see back label for listed weeds and control details.

<sup>^</sup>Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

<sup>‡</sup>Microtechnology signifies emulsion particles less than 1 micron.

**ACTIVE INGREDIENTS:**

2,4-D, dimethylamine salt*	0.885%
Mecoprop-p, dimethylamine salt**	0.573%
Dicamba, dimethylamine salt***	0.109%
OTHER INGREDIENTS <sup>††</sup> :	98.433%
	<b>Total 100.000%</b>
*2,4-D Acid Equivalent	0.738%
** Mecoprop-p Acid Equivalent	0.478%
*** Dicamba Acid Equivalent	0.091%

<sup>††</sup>Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN**

**Shake well before using**

**Protect from Freezing**

**See inside booklet for additional Precautionary Statements and Directions for Use**

**MADE IN CANADA (MAY CONTAIN IMPORTED PARTS) FOR:**

**Intelligro™, a Suncor Energy business**

2310 Lakeshore Road West

Mississauga, Ontario, Canada L5J 1K2

1-855-300-3997

EPA REG. NO. 69526-15  
641021

Patent(s): [www.suncor.com/patents](http://www.suncor.com/patents)  
 ™Trademark of Suncor Energy Inc. Used under license.

EPA EST. NO. 79693-CAN-001  
REV 021015

# PRODUCT SUMMARY

**CONTROLS:** Kills over 60 broadleaf weeds without harming lawns.^

^Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

**TREATMENT AREA :** 1 fl.oz. product treats up to 172 ft<sup>2</sup>

## MICROTECHNOLOGY :

Fast-acting, improved absorption formula

Patent-pending formulation improves herbicide absorption into weeds. Improved penetration reduces amount of active ingredients required in the formula to kill weeds.



# HOW IT WORKS

CIVITAS WEEDfree BRAND Concentrate♦ kills over 60 broadleaf weeds, without harming your lawn.^

Patent-pending formula improves the penetration of active ingredients into the weed.

♦see back label for listed weeds and control details

^Important: Do not apply to Bentgrass and Floratam. See full list of controlled weeds and use directions on peel away label.

## FIRST AID

### If swallowed:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

## HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. To reach a U.S. POISON CONTROL CENTER, call 1-800-222-1222. You may also contact INTELLIGRO, A SUNCOR ENERGY BUSINESS: 403-296-3000, or CANUTEC Canadian Transport Emergency Centre: 613-996-6666.

**NOTE TO PHYSICIAN:** May pose an aspiration pneumonia hazard, contains petroleum distillate.

# PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS.** Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

## Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are Barrier Laminate, Nitrile rubber, Butyl, Neoprene rubber or Viton. If you want more options, follow the instructions for Category E on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves (except for applicators using groundboom equipment), when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate, and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### **USER SAFETY RECOMMENDATIONS**

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking or ground water.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all label directions. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Read and follow all label directions.

Protect from freezing  
Shake well before using

### **NON-AGRICULTURAL USE REQUIREMENT**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.  
Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

## USE PRECAUTIONS



Do not apply as a fine mist because of potential injury to desirable plants. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

People or pets can enter treated area after sprays have dried.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, airblast, and chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles. When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for non-target species, non-target crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

#### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, beans, cotton, flowers, fruit trees (foliage), grapes (in growing stage), okra, ornamentals, soybeans (vegetative stage), sunflowers, tobacco, tomatoes, and other vegetables. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Do not spray exposed roots of ornamentals and trees.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for ground boom applications:

Do not apply with a nozzle height greater than 4 feet above the canopy.

**Precaution on warm season Grasses: Centipede, St. Augustine, Bahia, Common Bermuda, and Zoysia**

Some temporary discoloration may occur on warm season grasses. Do not apply if grass is stressed from heat or drought. Avoid application during transition periods from/to dormancy. **Centipede and St. Augustine** grasses are sensitive to 2,4-D and great care should be taken to prevent overdosing. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent discoloration or injury. Also, applications to dormant bermudagrass, dormant Zoysia grass, and dormant Bahiagrass are not suggested.

- **IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

**ADDITIONAL RESTRICTIONS**

- Do not apply through any type of irrigation system.
- Do not apply near lakes, streams, rivers ponds, or ditchbanks.
- Do not apply by air.
- Do not use more product than specified on the label.
- Do not apply when the lawn and/or turf, and/or site is under severe stress from drought, insect attack or heat. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not pour spray solutions near these plants. Do not apply in windy conditions.
- Do not irrigate within 24 hours of application.
- Do not tank mix with liquid fertilizers that contain nitrates.

<b>MAXIMUM APPLICATIONS PER YEAR</b>	<b>APPLICATION MAXIMUM</b>	<b>APPLICATION INTERVAL</b>
Limited to 2 applications per treatment area per year.	The maximum rate (8 gal./acre) for a single application is 0.75 lb ae/acre (0.9 lb DMA salt/acre) (0.424 lb 2,4-D ae, 0.274 lb MCP-p ae, 0.052 lb Dicamba ae per acre per application).	Repeat application if needed, but with no less than 30 days between applications.

# APPLICATION DIRECTIONS

## WEEDS CONTROLLED IN TURF or, LAWN AND ORNAMENTAL AREAS:

Bedstraw <sup>†</sup>	Grass leaved stitchwort <sup>†</sup>	Russian thistle <sup>†</sup>
Bindweed (field) <sup>†</sup>	Ground ivy (creeping Charlie) <sup>†</sup>	Sandwort (thyme-leaved) <sup>†</sup>
Birdsfoot trefoil <sup>†</sup>	Hawkweed <sup>†</sup>	Sheep sorrel <sup>†</sup>
Black medic	Heal-all <sup>†</sup>	Shepherd's purse
Blackseed plantain <sup>†</sup>	Henbit <sup>†</sup>	Smartweed (green) <sup>†</sup>
Blueweed <sup>†</sup>	Honeysuckle <sup>†</sup>	Soliva <sup>†</sup>
Brambles <sup>†</sup>	Jimsonweed <sup>†</sup>	Sow thistle <sup>†</sup>
Bull thistle <sup>†</sup>	Knotweed <sup>†</sup>	Speedwell (purslane leaved) <sup>†</sup>
Burdock <sup>†</sup>	Kochia <sup>†</sup>	Spotted spurge <sup>†</sup>
Buttercup <sup>†</sup>	Kudzu <sup>†</sup>	Spreadwell (annual) <sup>†</sup>
Canada thistle <sup>†</sup>	Lambsquarters <sup>†</sup>	Spurge <sup>†</sup>
Carolina geranium <sup>†</sup>	Little starwort <sup>†</sup>	Stitchwort <sup>†</sup>
Chickweed <sup>†</sup>	Mallow <sup>†</sup>	Stonecrop (mossy) <sup>†</sup>
Chicory <sup>†</sup>	Morningglory <sup>†</sup>	Sumac <sup>†</sup>
Clover (White clover in California)	Mouse-ear chickweed <sup>†</sup>	Veronica (thyme-leaved) <sup>†</sup>
Cocklebur <sup>†</sup>	Mustards <sup>†</sup>	Vervain <sup>†</sup>
Common chickweed	Narrow-leaved plantain	Vetch <sup>†</sup>
Cudweed <sup>†</sup>	Oxalis (yellow woodsorrel) <sup>†</sup>	Violet <sup>†</sup>
Daisy fleabane <sup>†</sup>	Ox-eye daisy <sup>†</sup>	Wild aster <sup>†</sup>
Dandelion	Pennywort <sup>†</sup>	Wild blackberry <sup>†</sup>
Devil's paint brush (hawkweed) <sup>†</sup>	Peppergrass <sup>†</sup>	Wild carrot <sup>†</sup>
Dock (curled) <sup>†</sup>	Pigweed <sup>†</sup>	Wild garlic <sup>†</sup>
English daisy <sup>†</sup>	Pineapple weed <sup>†</sup>	Wild geranium <sup>†</sup>
Evening primrose <sup>†</sup>	Plantains (narrow-leaved, buckhorn, broadleaf)	Wild lettuce <sup>†</sup>
Fall dandelion (fall hawkbit) <sup>†</sup>	Poison ivy <sup>†</sup>	Wild onion <sup>†</sup>
False dandelion <sup>†</sup>	Poorjoe <sup>†</sup>	Wild radish <sup>†</sup>
Fleabane <sup>†</sup>	Povertyweed <sup>†</sup>	Wild raspberry <sup>†</sup>
Florida pusley <sup>†</sup>	Prostrate pigweed <sup>†</sup>	Yarrow <sup>†</sup>
Frenchweed poison oak <sup>†</sup>	Purslane <sup>†</sup>	Yellow hawkweed <sup>†</sup>
	Ragweed <sup>†</sup>	

<sup>†</sup>Not registered for use in California

**TURF AND ORNAMENTAL (GOLF COURSES, CEMETERIES, PARKS,  
SPORT FIELDS, TURFGRASS, LAWNS AND OTHER GRASS AREAS)  
(Post emergence)**

**WHERE TO USE**

This product is for use as a broadcast or spot treatment on lawns for broadleaf weed control without damaging lawn grasses.

**IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass.

**NOTE:** Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

**WHEN TO USE**

- Do not water in 24 hours after spraying.
- Do not apply when rain or irrigation is anticipated.
- Spray when weeds are young and actively growing. Maximum control of weeds will be obtained from spring or early fall applications when weeds are actively growing.
- Do not water within 24 hours after spraying.
- Apply when daytime temperatures are above 45°F and below 85°F.
- Newly seeded lawns may be treated after the third mowing.
- If the lawn and/or turf is dry, water 24 hours before using this product.
- Grass seed can be sown 3 to 4 weeks after application.
- Do not apply when rain or irrigation is anticipated.
- Delay mowing 1 to 2 days before and after the application of this product.

**HOW TO USE**

**For broadcast treatment:**

Apply 2 to 8 gallons of the product per acre with the specified amount of water. Table below provides the recommended product rate and spray volume. Mix well before and during using. Use the lower rate (2-3 gal of the product/acre) on annual or small weeds. For best control of large weeds and tough perennial weeds, such as dandelion, use one application of the higher product rate ( $\geq$  4 gal of the product/acre), or two applications of lower rate making the second treatment 30 days following the first application.

<b>Product Rate (CIVITAS WEEDfree BRAND Concentrate<sup>†</sup>/acre)</b>	<b>Spray Volume (water/acre)</b>
2 gallons	10 to 15 gallons
3 gallons	15 to 20 gallons
4 gallons	20 to 50 gallons
5-6 gallons	50 to 80 gallons
6-8 gallons	60 to 120 gallons

<sup>†</sup>see back label for listed weeds and control details

Avoid overlapping between spray patterns.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

**For California, apply only 4-8 gallons/acre, as follows:**

4 gallons	20 to 50 gallons
5-6 gallons	50 to 80 gallons
6-8 gallons	60 to 120 gallons

**For spot treatment using hand-operated sprayers:**

Dilute 1 part of the product with 12 parts water to make about a 7.5% solution in water. Hold sprayer about 12 inches from weed and spray to lightly wet weeds.

**TURF, GROWN FOR SEED OR SOD (Post emergence)**

**WHERE TO USE**

For selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

**IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain grasses vary in tolerance to this product. Slight discoloration may occur, which will disappear after about one week.

**WHEN TO USE**

- Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 24 hours after application.
- Spray when weeds are young and actively growing. Apply when daytime temperatures are above 45°F and below 85°F.
- If the sod is dry, water 24 hours before using this product.
- Delay mowing 1 to 2 days before and after the application of this product.
- Treated areas may be reseeded 3 to 4 weeks after application.

**HOW TO USE**

**For broadcast treatment:**

Apply 2 to 8 gallons of the product per acre with the specified amount of water. Table below provides the recommended product rate and spray volume. Mixing well before and during use. Use the lower rate (2-3 gal of the product/acre) on annual or small weeds. For best control of large weeds and tough perennial weeds, such as dandelion, use one application of the higher product rate ( $\geq 4$  gal of the product/acre), or two applications of lower rate making the second treatment 30 days following the first application.

<b>CIVITAS WEEDfree BRAND Concentrate<sup>1</sup>/acre)</b>	<b>Spray Volume (water/acre)</b>
2 gallons	10 to 15 gallons
3 gallons	15 to 20 gallons
4 gallons	20 to 50 gallons
5-6 gallons	50 to 80 gallons
6-8 gallons	60 to 120 gallons

<sup>1</sup>see back label for listed weeds and control details

Avoid overlapping between spray patterns.



Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

**For California, apply only 4-8 gallons/acre, as follows:**

4 gallons	20 to 50 gallons
5-6 gallons	50 to 80 gallons
6-8 gallons	60 to 120 gallons

**For spot treatment using hand-operated sprayers:**

Dilute 1 part of the product with 12 parts water to make about a 7.5% solution in water. Hold sprayer about 12 inches from weed and spray to lightly wet weeds.

**NEWLY SEEDED AREAS:** Applying this product to grass seedlings is recommended after the third mowing.

**NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS:** Delay applications to newly sodded, sprigged, or plugged grasses 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant Bermudagrass, dormant Zoysia grass, and dormant Bahiagrass are not suggested.

## EXPECTED RESULTS



Visible injury on weeds can be seen within 12 hours. It can take up to 4 weeks for weeds to completely die.

How often to apply:

Repeat application can be made after 30 days if necessary. The maximum number of broadcast applications per treatment site is 2 per year.

## STORAGE AND DISPOSAL



Do not contaminate water, food, or feed by storage and disposal.

### PESTICIDE STORAGE:

Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place in a locked storage area. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons.

Confine spills. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

### PESTICIDE DISPOSAL:

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

### CONTAINER HANDLING:

#### ***Rigid, nonrefillable containers, equal to or less than five gallons:***

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

***Rigid nonrefillable containers, greater than 5 gallons:***

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

***Non-rigid, nonrefillable container (275-gallon bladder with frame):***

Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling is unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Return frame to dealer.

## **CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY**

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Intelligro, a Suncor Energy business, the manufacturer or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law neither Intelligro, a Suncor Energy business, the manufacturer, nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product. To the extent consistent with applicable law or except as warranted by this label, Intelligro, a Suncor Energy business, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and Intelligro, a Suncor Energy business, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.

4.75"

6.50"

7.25"

5.75"

Folded Onsert

Maximum size area

for label graphics (under Onsert)

**ACTIVE INGREDIENTS:**

2,4-D Acid Equivalent	0.885%
Acifluorfen	0.000%
Mecoprop-p, dimethylamine salt**	0.573%
Dicamba, dimethylamine salt***	0.109%
OTHER INGREDIENTS††	98.433%
<b>Total</b>	<b>100.000%</b>

\*2,4-D Acid Equivalent  
\*\* Mecoprop-p Acid Equivalent  
\*\*\* Dicamba Acid Equivalent  
†† Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN**

EPA REG. NO. 68926-15  
EPA EST. NO. 79693-CALH-001  
Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.  
\*See back label for listed weeds and control details.  
Patent(s): [www.suncoor.com/patents](http://www.suncoor.com/patents)

See inside booklet for additional Precautionary Statements and Directions for Use  
**MADE IN CANADA (MAY CONTAIN IMPORTED PARTS) FOR:**  
Intelligro, a Suncoor Energy business  
2310 Lakeshore Road West, Mississauga, Ontario, Canada L5J 1K2  
1-855-300-5987

<b>FIRST AID</b>
• Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
<b>HOTLINE NUMBER</b>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. To reach a U.S. POISON CONTROL CENTER, call 1-800-222-1222. You may also contact INTELLIGRO, A SUNCOOR ENERGY BUSINESS: 403-293-5900 or CANULTEC Canadian Transport Emergency Centre: 613-985-5966.
<b>NOTE TO PHYSICIAN:</b> May pose an aspiration pneumonia hazard, contains petroleum distillate.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

**STORAGE AND DISPOSAL**

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage and disposal.  
**PESTICIDE STORAGE:** Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place, preferably in a locked storage area.  
**PESTICIDE AND CONTAINER HANDLING:** Wash hands thoroughly with soap and water after handling. Do not eat, drink, or use tobacco while handling. Offer for recycling if available, or reuse container in trash.  
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.



**CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY**

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other material, or the manner of use or application, all of which are beyond the control of Intelligro, a Suncoor Energy business, the manufacturer or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law neither Intelligro, a Suncoor Energy business, the manufacturer or seller, nor the manufacturer or seller shall be liable for consequential damages. Intelligro, a Suncoor Energy business, the manufacturer or seller, makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or applicability of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and Intelligro, a Suncoor Energy business, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.

641021

**NET CONTENTS:**  2.5 U.S. GAL (9.5L)  2x2.5 U.S. GAL (2x9.5L)  
 30 U.S. GAL (113.5L)  275 U.S. GAL (1040L)



DESIGN LIMITED  
BEI SANSON, CRESCENT  
176 THE BRIDGE  
SEAPORT TOWNSHIP, ONTARIO  
L7G 1H2  
TEL: (905) 877-5122  
FAX: (905) 877-5122

CIVITAS WEED FREE BRAND (US)	PROGRAM: ADOBE ILLUSTRATOR CC
PROJECT DATE: DEC. 16/2014	PRINTED ON:
DOCKET NO.: M-121	COLOURS: BLACK
SCALE: DISTORTED: <input type="checkbox"/> YES <input type="checkbox"/> NO	FONTS:
LABEL NO.: 641021 (THIS IS A NEW SKU)	
LAST REVISION DATE: DEC 18/2014 (COPY REVISIONS), DEC 19/2014 (COPY REVISIONS), JAN 15/2015 (COPY REVISIONS), JAN 23/2015 COPY REVISIONS, JAN 26/2015 (COPY REVISIONS), FEB. 2/2015 (COPY REVISIONS), FEB. 10/2015 (COPY REVISIONS).	